I. Citation or Source:		
Banks, R.C. 1979. Human related mortality of birds in the United States. U.S. Fish & Wildlife Service, National Fish and Wildlife Lab, Special Scientific Report – Wildlife No. 215:1-16. GPO 848-972.		
Source Type (check one): Mortality Summary		
Peer-reviewed Paper √ Other (specify):  Agency Report  Conference Proceedings		
II. Objectives (list)		
Summarize avian mortalities from different sources. Banks indicates the estimate of avian kills under "normal conditions." Estimates 2,500 bird Mountains. If only half of towers presents hazards, 1979 estimates 1,25	kills/tower/year. No mass mortalities in o	
III. Species		
Addressed mortality factors by family or groups.		
IV. Study Methods (briefly list)		
N/A		
V. Duration of Study N/A		
Duration (provide dates): Single Year	Seasons: Spring Migration Both	·
Multiple Years	Fall Migration Year	long

VI. Carcass Search Methods (if applicable) N/A
Search Conditions: Daily Weekly Only after overcast nights with a low celling or storm events
Other Periods (Describe):
Search Blases Evaluated, Including Observer Blas and Scavenger Activity? Yes No √
Search Area Described? Yes No √
VII. Analytical and Statistical Methods N/A
Statistical method(s) used: (list)
N/A
Comments:
VIII. Number of Tower Sites: N/A Proximity:
VIII. Number of Tower Sites: N/A Proximity:
IX. Behavioral Observations at the Tower: Yes No √ Describe if applicable to statement or conclusion being evaluated.
Describe it applicable to statement or conclusion being evaluated.
V. Danimantation of Worthon Featons? Von No.s/
X. Documentation of Weather Factors? Yes No √ Describe if applicable to statement or conclusion being evaluated.

XI. Inclusion of Structural and Landscape Conditions? Yes No √ Describe if applicable to statement or conclusion being evaluated.
XII. Brief Description of Results
Discusses overall bird mortality reports at communication towers to date. Most reports were of mass mortalities, which is of limited value in establishing annual mortality rates under "normal conditions." Estimates if only 50% of towers at that time presented an avian collision risk, an estimated 2.500 birds per tower could be affected, totaling an estimated 1,250,000 birds per year in the U.S. Vireonidae, Parulidae, and Fringillidae are the most frequently affected bird families.
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)
Are additional studies identified? Yes No √ If yes explain and list studies.
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.

I. Citation or Source:	
Ball, L.G., K. Zyskowski, and G. Escalona-Segura. 1995. Recent bird to Society Bulletin 46(4):33-36.	nortality at a Topeka television tower. Kansas Ornithological
Source Type (check one): Incident report	
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings	
II. Study Objectives (list)	
Monitoring number of bird kills and species recorded.	
III. Species	
Detailed species list: 2,808 bird mortalities, 91 species. Gray catbird an most common for October dates. A number of larger, water birds, wren other passerines recorded.	
IV. Study Methods (briefly list)	
Focus was to document otherwise rare occurrences of birds in area.	
V. Duration of Study 4 events	
Duration (provide dates):  Single Year  Multiple Years 25-26 Sep 1985, 30 Sep - 1 Oct 1986,  11-12 Oct 1986, 8-9 Oct 1994.	Seasons: Spring Migration Both Fall Migration ✓ Yearlong

VI. Carcass Search Methods (if applicable)
Search Conditions: Daily Weekly Only after overcast nights with a low ceiling or storm events
Other Periods (Describe): Based on events (four total).
Search Biases Evaluated, Including Observer Bias and Scavenger Activity? Yes No √
Search Area Described? Yes No √
VII. Analytical and Statistical Methods N/A
Statistical method(s) used: (list)
Summer mentody) used. (use)
Comments:
VIII. Number of Tower Sites:   Proximity:
KTKA tower 4 km west of Topeka, Kansas.
439 m tall. Guyed. Incandescent (red?) lights.
IX. Behavioral Observations at the Tower: Yes No ↓  Describe if applicable to statement or conclusion being evaluated.
X. Documentation of Weather Factors? Yes √ No  Describe if applicable to statement or conclusion being evaluated.
Low cloud ceiling following major decrease in wind speed and preceding temperatures falling. Only precipitation occurred 11-12 Oct 1986, which coincided w/ rain and light snow.

XI. Inclusion of Structural and Landscape Conditions? Yes No √ Describe if applicable to statement or conclusion being evaluated.
XII. Brief Description of Results
No scientific data, but good information on potential effects to rare, threatened, or endangered species. Detailed species list.
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)
Are additional studies identified? Yes No √ If yes explain and list studies.
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.

L Citation or Source:		
Avery, M.L., P.F. Springer, and J.F. Cassel. 1978. The composition and seasonal variation of bird losses at a tall tower in southeastern North Dakota. American Birds 32(6):1141-1121.		
Source Type (check one): Study		
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings		
II. Study Objectives (list)		
Determine extent and seasonal variation of bird mortality.		ĺ
III. Species		
937 (partial sample) (1,075 mean annual) 102 species of birds; 46% neotropical migrants; predominantly vireos a 4, 298 estimated for 5 seasons 1971-73.	and warblers.	
IV. Study Methods (briefly list)		
Surveyed at dawn almost daily. Strata w/in 151 ft surveyed 100%. Randomized sampling effort for 3 other concentric strata (302, 600, and Total 8 sampling plots for each concentric circle (40.7 ft ea. side ea. ple Used nets on sampling plots.		
V. Duration of Study: 3 years - nightly		
Duration (provide dates):	Seasons:	
Single Year  Multiple Years 1971-1973 (3 yrs)  Except for 7 days: daily @ dawn  30 Mar - 4 Jun (1972) 2 Apr - 2 Jun (1973)  8 Aug - 15 Nov (1972) 12 Aug - 3 Nov (1973)  (surveyed other adjacent days)	Spring Migration	Both √ Yearlong

VI. Carcass Search Methods (if applicable)
Search Conditions: Daily \( \frac{1}{2} \) Weekly Only after overcast nights with a low ceiling or storm events
Other Periods (Describe):
Search Biases Evaluated, Including Observer Bias and Scavenger Activity? Yes √ No Scavenger Study
Search Area Described? Yes √ No
Scavenger Study: planted 296 birds at various points over 4 seasons Results: Range 2.4% carcass removal in spring 1972 (7.3% nightly); 17.6% carcass removal in spring 1973. Removal averaged 7.4% overall.
Marshy/grass)and search area.
VII. Analytical and Statistical Methods
Statistical method(s) used: (list)
Predominantly descriptive study. Chi-square tests used to determine if kill rates of particular species or families varied w/ season.
Comments:
VIII. Number of Tower Sites: 1 Proximity:
U.S. Coast guard Omega Navigational Station – James River Valley; 3 km W. LaMoure, ND. 5 red, non-flashing, obstruction lights
4 red, flashing 700-W beacons 366-meter tower; guyed; 16 evenly spaced transmitting cables from top of antennae to a perimeter road 732 from tower.
200 11000 100101, 50-7011, 10-1
IX. Behavioral Observations at the Tower: Yes No \( \square \)
Describe if applicable to statement or conclusion being evaluated.
X. Documentation of Weather Factors? Yes No √ Describe if applicable to statement or conclusion being evaluated.
l l

XI. Inclusion of Structural and Landscape Conditions? Yes No √ Describe if applicable to statement or conclusion being evaluated.
XII. Brief Description of Results
Consistent sampling effort; measured scavenger removal estimated @ 7.3% nightly (7.4% overall).  54% kills occurred in fall migration (Sep-Nov) = Stratum D (183-732 m from tower) suggest that most mortality caused by guy wires/transmitting cables farther from towers.  44% warblers and 11% vireos over half of mortalities recorded primarily in fall.  Fall mortalities typically not local breeders.  Wrens, icterids, and fringillids predominantly spring kills.  Spring mortalities commonly local breeding birds, inferring that local breeders more affected than migrants heading north. Hypothesize local breeders more selective in spring as compared to fall as to where perch after nights migration. Birds breed in habitats similar to Omega station are more prone to collisions. Also, diurnal collisions w/ local breeders may occur, particularly during inclement weather or poor visibility. Seasonal variation in species composition.
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)
Are additional studies identified? Yes No ⊻ If yes explain and list studies.
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.

I. Citation or Source:		
Avery, M.L., P.F. Springer, and J.F. Cassel. 1977. Weather influences on nocturnal bird mortality at a North Dakota tower. Wilson Bulletin 89(2):291-299.		
Source Type (check one): Study		
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings		
II. Study Objectives (list)		
Record weather influences on bird mortalities.		i
III. Species		
~ 1, 064 birds collected.		
IV. Study Methods (briefly list)		
4 concentric strata (strata A = 100%; other strata lower percentage cove	er).	
V. Duration of Study: 4 migrational seasons		
Duration (provide dates):	Seasons:	
Single Year  Multiple Years 1972-1973 (2 yrs)  30 Mar - 4 Jun (1972) 2 Apr - 2 Jun (1973)  8 Aug - 15 Nov (1972) 12 Aug - 3 Nov (1973)  nightly searches @ daybreak (except 7 days)	Spring Migration	Both √ Yearlong

VI. Carcass Search Methods (if applicable)
Search Conditions: Daily √ Weekly Only after overcast nights with a low celling or storm events
Other Periods (Describe):
Search Biases Evaluated, Including Observer Bias and Scavenger Activity? Yes No √
Search Area Described? Yes √ No
VII. Analytical and Statistical Methods
Statistical method(s) used: (list)
<ul> <li>Chi-square-goodness-of-fit test – w/in ea family; losses in entire peak periods were same proportion to number of nights in those categories.</li> <li>G-test – determine independence b/w cloud cover and distance of kill and b/w cloud cover and season.</li> </ul>
Comments:
VIII. Number of Tower Sites: 1 Proximity:
U.S. Coast guard Omega Navigational Station – James River Valley; 3 km W. LaMoure, ND.  5 red, non-flashing, obstruction lights  4 red, flashing 700-W beacons  366-meter tower; guyed; 16 evenly spaced transmitting cables from top of antennae to a perimeter road 732 from tower.  Omega tower unique because of 16 transmitting cables - increase collision risk, particularly farther from base on clear nights.
IX. Behavioral Observations at the Tower: Yes √ No Describe if applicable to statement or conclusion being evaluated.
Similar to Cochran and Graber 1958 and Avery et al., 1976. Mortality information suggests behavioral differences that depict where families or group-level birds may be affected.
X. Documentation of Weather Factors? Yes √ No Describe if applicable to statement or conclusion being evaluated.
Hourly weather reports from FAA Flight Service Station @ Jamestown (72 km N-NW of LaMoure).  Precipitation; wind speed and direction recorded.  On overcast nights – losses concentrated near the tower (and lights) in strata A and B; non-overcast nights – more evenly distributed.  Spring and fall – difference in mortality b/w overcast and non-overcast was statistically significant. Infers distance of losses from tower was influenced by cloud cover.

XI. Inclusion of Structural and Landscape Conditions? Yes \( \sqrt{No} \)  Describe if applicable to statement or conclusion being evaluated.	
Marshy/grassland.	
XII. Brief Description of Results  Consistent sampling effort, major fall kills followed cold fronts.	
Table 1: 5 largest single-night losses in 1972-73; 4 of 5 nights overcast. Fall losses on overcast nights occurred w/in 12 hours of cold fronts, which is consistent w/ other studies (as listed). Spring kills more evenly distributed; no direct association w/ frontal movements; majority of kills occurred w/ favorable (SE) winds.  Ceilometer observations – majority of spring migration occurred w/ SE winds.	
The percent of fall mortalities were higher than the percent recorded in the spring migration w/in 92 m of the tower; beyond 92 m this is reversed except in 1972. In summary, larger spring losses consistently occurred at greater distances from the tower than in the fall.	
16 transmitting cables increase the collision risk. During clear (non-overcast) nights, birds avoided towers, but higher collisions recorded farther from tower.	
Mortality data infer/suggest difference in bird strikes (number, location, weather conditions) relates to family or bird group.	
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)	
Are additional studies identified? Yes √ No If yes explain and list studies.	
Differences among various taxa of nocturnal migrants in responses to tall, lighted structures warrants further research. Conceivably, data may provide methods where losses to same species at towers could be reduced.	
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)	
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.	

I. Citation or Source:		
Avery, M.L., P.F. Springer, and J.F. Cassel. 1976. The effects of a ta study. Auk 93(2):281-291.	ll tower on nocturnal bird migration	- a portable ceilometer
Source Type (check one): Study		
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings		
II. Study Objectives (list)		
Record bird behavior at and away from tower. Examine bird distribution @ towers. Monitor numbers of birds at and away from tower. Record bird behavior near tower lighting. Monitor movements.		
III. Species		
Three species frequently recorded at the tower: sora, common yellow	vthroat, and savannah sparrow.	
IV. Study Methods (briefly list)		
Nighttime observations of migrants using portable ceilometer techniq Monitored/recorded weather conditions. Used binoculars and spotting scope.	jue.	
V. Duration of Study 4 nights/week		
Duration (provide dates):	Seasons:	
Single Year	Spring Migration	Both √
Multiple Years 1972, 1973  18 April – 1 June (1972)  2 Apr – 31 May (1973)  19 Apr. – 26 Oct (1972)  16 Apr. – 27 Oct (1973)	Fall Migration	Yearlong

VI. Carcass Search Methods (if applicable)
Search Conditions: Daily Weekly Only after overcast nights with a low ceiling or storm events
Other Periods (Describe): 4 nights/week (see V.)
Search Biases Evaluated, Including Observer Bias and Scavenger Activity? Yes No √
Search Area Described? Yes 🗸 No
VII. Analytical and Statistical Methods
Statistical method(s) used: (list)
N/A
Comments:
VIII. Number of Tower Sites:   Proximity: N/A
U.S. Coast guard Omega Navigational Station – James River Valley; 3 km W. LaMoure, ND. 5 red, non-flashing, obstruction lights
4 red, flashing 700-W beacons
366-meter tower; guyed; 16 evenly spaced transmitting cables from top of antennae to a perimeter road 732 from tower.
IX. Behavioral Observations at the Tower: Yes \( \subset \) No
Describe if applicable to statement or conclusion being evaluated.
"Fluttered/milled about" oriented mainly into wind (did <u>not</u> orient toward tower lights). Flight pattern: several wingbeats – brief
pause – (whether transmitting or not). Foggy @ dawn 26 Aug 1973; 35 m near red tower lights – flew upwind, frequently
pausing/fluttering ~20 m SE of tower – turn slightly – blown downwind ~ 50 m NW of tower – stopped and began slow flight upwind again. Counterclockwise – narrow elliptical path.
V. Dooumantation of Weather Feetage? Von./ No.
X. Documentation of Weather Factors? Yes √ No Describe if applicable to statement or conclusion being evaluated.
Data grouped by season (spring or fall).
Within each seasonal group, sightings divided into overcast or nonovercast (clear or partly cloudy) classes.  Classes subdivided by location (at tower or 305 m to NE of tower).

XI. Inclusion of Structural and Landscape Conditions? Yes No √ Describe if applicable to statement or conclusion being evaluated.
XII. Brief Description of Results
Overcast nights each season - number of migrants observed @ tower significantly greater than number observed 305 m to NE of tower. Clear nights - reverse was true. Partly cloudy nights - not significant in fall. Directional movements under various conditions; multiple seasons recorded. Overcast nights w/ clearing showed number birds sharply decreasing @ tower. Number of birds decreased/increased when lights turned off/on 22-23 Aug 1973. Inference: congregation of nocturnal migrants - orientation using celestial cues. More likely: reluctant to leave area of illumination when passing tower, particularly during inclement weather. Migrants may actively avoid tower on clear nights.
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)
Are additional studies identified? Yes No √ If yes explain and list studies.
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.

I. Citation or Source:		
Crawford, R.L. and R.T. Engstrom. 2001. Characteristics of avian mo Field Ornithol. 72(3):380-388.	rtality at a north Florida television to	ower: A 29-year study. J.
Source Type (check one): Study		
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings		
II. Study Objectives (list)		
Determine extent of avian mortality over long study period.  1) Summarize 29 years of data.  2) Examine effects of tower height on avian mortality.  3) Evaluate effects of scavengers on number of mortalities determined.	octed.	
III. Species		
44,007 bird kills of 186 species over 29-year period.  > 94% were neotropical migrants; red-eyed vireo #1 mortality recorded After tower shortened, only 32 bird mortalities recorded Oct 1999 (27 v Of 41 families; Parulidae and Vireonidae = 64%; primarily neotropical 99% of mortality concentrated in only 15 of the 43 total families record	risits); 14 recorded Oct 2000 (18 vis migrants/noctumal migrants.	its)
IV. Study Methods (briefly list)		
Attempted daily surveys at dawn. Descriptive study; information not quantified.		·
V. Duration of Study: 29 years		<del></del>
Duration (provide dates):  Single Year  Multiple Years 1955-1967-1983 (29 years)  Main part of study concluded 1985. 284-ft tower also checked in Oct 1999 and 2000 for comparison w/ taller tower for same period.	Seasons: Spring Migration Fall Migration	Both √ Yearlong

VI. Carcass Search Methods (if applicable)
Search Conditions: Daily √ Weekly Only after overcast nights with a low ceiling or storm events
Other Periods (Describe):
Search Biases Evaluated, Including Observer Bias and Scavenger Activity? Yes 🗹 No
Search Area Described? Yes √ No
20-acre mowed area under tower and guy wires.  Aggressive scavenger removal using live trapping and poisons from 1955-1967 (13 years) and again 1974-1976 (3 years).  Determined strong scavenger effect; reduced predator control resulted in 71% reduction in birds found.
VII. Analytical and Statistical Methods Largely descriptive
Statistical method(s) used: (list)
Used ANOVA to determine if kill rates changed when tower height increased from 669 to 1,010 feet. No effect reported.
Comments:
VIII. Number of Tower Sites: 1 Proximity:  WCTV tower northern Leon co, FL (Tall Timbers Research Station reporting)  1955-1960 = 669 feet  1960 = new tower replaced 1,010 feet  1989 = tower shortened to 295 feet
IX. Behavioral Observations at the Tower: Yes No √ Describe if applicable to statement or conclusion being evaluated,
X. Documentation of Weather Factors? Yes No Describe if applicable to statement or conclusion being evaluated.
Crawford and Engstrom controlled for weather and scavenger conditions.

XI. Inclusion of Structural and Landscape Conditions? Yes No √ Describe if applicable to statement or conclusion being evaluated.
XII. Brief Description of Results
Examines relationship of tower height (3 towers at same site) to bird kills. Authors state that towers less than 295 feet may not present serious risk for bird collisions. Scavenger control recommended.
Mean mortality numbers = 1,517/year overall. With scavenger control the mean = 2,248 mortalities/year; without scavenger control the mean = 642.
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)
Are additional studies identified? Yes No √ If yes explain and list studies.
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.

I. Citation or Source;		
Crawford, R.L. 1971. Predation on birds killed at TV tower. Oriole 36:	33-35.	
Source Type (check one): Study		
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings		
II. Study Objectives (list)		
Descriptive summary of predation issues on bird mortalities at tower sit	es.	
III. Species		
N/A		.1
IV. Study Methods (briefly list)		
Tall Timber Bulletin Number 1 (1962) and Number 8 (1967) have detain Planted birds on tower site for 5 nights to monitor/calculate scavenging		966.
V. Duration of Study		
Duration (provide dates):	Seasons:	
Single Year <u>5 nights 21-27 Oct 1971</u> Multiple Years	Spring Migration Fall Migration √	Both

VI. Carcass Search Methods (if applicable)
Search Conditions: Daily Weekly Only after overcast nights with a low ceiling or storm events √
Other Periods (Describe):
Search Biases Evaluated, Including Observer Bias and Scavenger Activity? Yes V No Scavenger Removal
Search Area Described? Yes No 🗹
1955-1966/ 1971 5 nights = scavenger removal monitored.
VII. Analytical and Statistical Methods
Statistical method(s) used: (list)
N/A
Comments:
VIII. Number of Tower Sites: 1 Proximity:
WCTV tower northern Leon co, FL (Tall Timbers Research Station reporting) 1955-1960 = 669 feet 1960 = new tower replaced 1,010 feet
IX. Behavioral Observations at the Tower: Yes No √ Describe if applicable to statement or conclusion being evaluated.
X. Documentation of Weather Factors? Yes No √ Describe if applicable to statement or conclusion being evaluated.

XI. Inclusion of Structural and Landscape Conditions? Yes No √ Describe if applicable to statement or conclusion being evaluated.
XII. Brief Description of Results
Predation was high. Total predation rates (day and night) were 93% (147 of 157 birds). Great homed owls were the primary nocturnal scavengers, with common and fish crows scavenging diurnally.
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)
Are additional studies identified? Yes ⊻ No If yes explain and list studies.
Tower studies should account for predation and scavenger removal rates. Recommended strict predator control, early morning surveys, and whether drastic predator control measures are worth the data.
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.

· · · · · · · · · · · · · · · · · · ·		
I. Citation or Source:		
Crawford, R.L. 1978. Autumn bird casualties at a northern Florida TV	Fower: 1973-1975. Wilson Bulletin 90(3):335-345.	
Source Type (check one): Study		
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings		
II. Study Objectives (list)		
Record age and sex data for 3, 223 birds killed during fall periods (Aug-Nov) of 1973-1975.		
III. Species		
3,864 bird mortalities reported; 109 species  Mass Mortalities: 17 Oct 1973 = (133); 5 Sep 1974 (134); 23 Sep 1974 (220); 17 Oct 1974 (971); 14 Sep 1975 (636); 15 Sep 1975 (486)  Aug-Nov only: 1973 = 261 birds; 57 species 1974 = 1,832 birds, 87 species (w/ predator control) 1975 = 1,771 birds; 90 species (w/ predator control)		
IV. Study Methods (briefly list)		
Almost daily searches. Reinstituted rigorous predator control in 1974-75, which explains increased carcass retrieval rates. Analyzed sex and age ratios.		
V. Duration of Study 3 years		
Duration (provide dates): Single Year Multiple Years 1973-1975	Seasons:  Spring Migration Both  Fall Migration (Aug – Nov) \( \frac{1}{2} \) Yearlong	

VI. Carcass Search Methods (if applicable)
Search Conditions: Daily √ Weekly Only after overcast nights with a low ceiling or storm events
Other Periods (Describe):
Search Biases Evaluated, Including Observer Bias and Scavenger Activity? Yes 🗸 No
Search Area Described? Yes No ⊻
Scavenger/predator removal program in place.
VII. Analytical and Statistical Methods
Statistical method(s) used: (list)
Variance test for homogeneity of the binomial distribution.
Comments:
VIII. Number of Tower Sites: 1-2 Proximity:
WCTV – Leon Co. Florida 1,008-foot, guyed TV tower (see Crawford and Engstrom 2001). Steady and flashing red lights. WCTV tower was focus, but these tower results compared to central peninsular tower (WDBO) by Taylor and Anderson, 1973.
IX. Behavioral Observations at the Tower: Yes No √ Describe if applicable to statement or conclusion being evaluated.
X. Documentation of Weather Factors? Yes \( \sqrt{No} \)  Describe if applicable to statement or conclusion being evaluated.
Most mortalities occurred following cold fronts.

XI. Inclusion of Structural and Landscape Conditions? Yes No √ Describe if applicable to statement or conclusion being evaluated.
XII. Brief Description of Results
Compared different types of species' migration. Authors state that differences in migration movement periods result in differences in mortality events (e.g., displaced female ruby-crowned kinglets).  Larger number of adults (earlier migrants) than immature birds recorded (Taylor and Anderson, 1973).  The two tower comparisons and sampling delineated different migration systems.  Strong inference to mass mortalities associated w/ cold fronts.  Suggests those species that migrate early are less likely to be impacted by cold fronts and hence collisions.  Also see Nolan and Mumford 1965.
XIII. Need for and Scope of Additional Studies (Only applicable if new data or study is provided.)
Are additional studies identified? Yes No √ If yes explain and list studies.
XIV. Suggested Methods to Minimize Impacts (Only applicable if new data or study is provided.)
Are specific methods identified? Yes No √ If yes explain and list specific mitigative methods.

I. Citation or Source:		
Stoddard, H.L., Sr. 1962. Bird casualties at a Leon County, Florida TV tower: 1955-1961. Bull. Tall Timbers Res. Sta. 1:94.		
Source Type (check one): Study.		
Peer-reviewed Paper √ Other (specify): Agency Report Conference Proceedings		
II. Study Objectives (list)		
Initiate a long-term study of bird collisions w/ towers. Record species and number of birds killed. Document extent of scavenger removal of carcasses.		
III. Species		
15, 251 total birds; 149 species See page 3 of report for listings of mass mortality kills. Few to several species of bats recorded.		
IV. Study Methods (briefly list)		
Almost daily surveys.		
V. Duration of Study 7 years		
Duration (provide dates): Single Year Multiple Years 1955-1961	Seasons: Spring Migration Fall Migration	Both √ Yearlong